

An X-ray of your shoulder; what happens next?

You have been referred to the Radiology department. An X-ray image will be made to see if you have a fracture.

If you have a fracture

If the X-ray shows you have a fracture, you will be sent to the emergency room for treatment. You can report at the reception desk. The receptionist will tell you what will happen next.

If you don't have a fracture

If the X-ray doesn't show a fracture, it means your shoulder is merely bruised. No further treatment will be necessary. You are free to go home. The physician will examine your X-ray again the next day. They will give you a call if necessary. We will inform your own general practitioner that you visited the out-of-hours medical centre. They will also receive the results of your X-ray exam.

Self-care:

- Keep your shoulder cool. Use an ice pack or something else from your freezer. Wrap this in a washcloth or tea towel and place it on your shoulder for 10 minutes. Do this several times a day for two days.
- Move your shoulder carefully. On the first day, you can start with some hand exercises. Open and close your hand to make fist. This exercise keeps the muscles in your arm active. Stop moving if it hurts too much.
- If you want, you can wear a sling. Make sure your upper arm hangs against your body in a relaxed manner. You can buy a sling at your local pharmacy or chemist's. Scan the QR code (at the bottom right of this page) for an instructional video. You can also make an appointment with your physical therapist to work on your recovery.

Pain medication (adults):

- You can take paracetamol for the pain: two 500-mg tablets of paracetamol 3 to 4 times a day.
- If paracetamol is not enough to relieve your pain, you can also take ibuprofen: one 400-mg tablet 3 times a day. Note: only take ibuprofen if you are sure you can do so safely. If in doubt, consult your physician or pharmacist.

Pain medication (children):

- You can give your child paracetamol: check the packaging for the recommended dose.
- If paracetamol is not enough to relieve your child's pain, you can also give them liquid Nurofen. Check the packaging for the recommended dose.
- Children aged 12 or over may also take ibuprofen: check the packaging for the recommended dose.

Contact your own general practitioner if:

- the above recommendations are not effective
- your complaints persist for more than 5-7 days
- you are worried
- you have any questions

